

Table 1

Recommended Child and Adolescent Immunization Schedule for Ages 18 Years or Younger, United States, 2025

These recommendations must be read with the notes that follow. For those who fall behind or start late, provide catch-up vaccination at the earliest opportunity as indicated by the green bars. To determine minimum intervals between doses, see the catch-up schedule (Table 2).

Vaccine and other immunizing agents	Birth	1 mo	2 mos	4 mos	6 mos	9 mos	12 mos	15 mos	18 mos	19–23 mos	2–3 yrs	4–6 yrs	7–10 yrs	11–12 yrs	13–15 yrs	16 yrs	17–18 yrs
Respiratory syncytial virus (RSV-mAb [Nirsevimab])	1st dose	1 dose depending on maternal RSV vaccination status (See Notes)								1 dose (8 through 19 months); See Notes							
Hepatitis B (HepB)	1st dose	← 2nd dose →			← 3rd dose →												
Rotavirus (RV): RV1 (2-dose series), RV5 (3-dose series)			1st dose	2nd dose	See Notes												
Diphtheria, tetanus, acellular pertussis (DTaP <7 yrs)			1st dose	2nd dose	3rd dose			← 4th dose →				5th dose					
<i>Haemophilus influenzae</i> type b (Hib)			1st dose	2nd dose	See Notes		← 3rd or 4th dose (See Notes) →										
Pneumococcal conjugate (PCV15, PCV20)			1st dose	2nd dose	3rd dose		← 4th dose →										
Inactivated poliovirus (IPV)			1st dose	2nd dose	3rd dose		← 3rd dose →					4th dose					See Note
COVID-19 (1vCOV-mRNA, 1vCOV-aPS)									1 or more doses of 2024–2025 vaccine (See Notes)								
Influenza (IIV3, cclIV3)									1 or 2 doses annually					1 dose annually			
Influenza (LAIV3)													1 or 2 doses annually	or 1 dose annually			
Measles, mumps, rubella (MMR)					See Notes		← 1st dose →					2nd dose					
Varicella (VAR)							← 1st dose →					2nd dose					
Hepatitis A (HepA)					See Notes			2-dose series (See Notes)									
Tetanus, diphtheria, acellular pertussis (Tdap ≥7 yrs)																1 dose	
Human papillomavirus (HPV)																See Notes	
Meningococcal (MenACWY-CRM ≥2 mos, MenACWY-TT ≥2 years)								See Notes							1st dose		2nd dose
Meningococcal B (MenB-4C, MenB-FHbp)																	
Respiratory syncytial virus vaccine (RSV [Abrysvo])																	
Dengue (DEN4CYD: 9–16 yrs)																	Seasonal administration during pregnancy (See Notes)
Mpox																	Seropositive in endemic dengue areas (See Notes)

Range of recommended ages for all children
 Range of recommended ages for catch-up vaccination
 Range of recommended ages for certain high-risk groups or nonimmunizations
 Recommended vaccination can begin in this age group
 Recommended vaccination based on shared clinical decision-making
 No Guidance/Not Applicable